



# Statistical Control Techniques: Partial and Semipartial Correlation

## Chapter 7



# Today's Lecture

- Statistical control techniques.
- Partial correlation.
- Semipartial correlation.

Overview

● Today's Lecture

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Statistical Control

## Overview

### Statistical Control

- Visualizing Control
- Example Data Set

## Partial Correlation

## Measurement Errors

## Semipartial Correlation

## Multiple Partial Correlation


## Wrapping Up

- Statistical control can be equated to controlling for variability.
- In experimental research (with human subjects), this is often accomplished by random assignment to a set of experimenter controlled conditions.
- In quasi-experimental research random assignment cannot be accomplished, although experimental conditions may exist.
- In nonexperimental or observational research, the “experimenter” does not control anything...but can observe many things.
- To obtain control when none is present, statistics can be useful.



# Venn Diagrams

One way to visualize the concepts described in this lecture is to imagine the variability of a variable as being represented by a circle.



Overview

Statistical Control

● Visualizing  
Control

● Example Data Set

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation


Multiple Partial  
Correlation

Wrapping Up



# Venn Diagrams

Using  $X_1$  to predict  $Y$ :



Overview

Statistical Control

● Visualizing  
Control

● Example Data Set

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation


Multiple Partial  
Correlation

Wrapping Up



# Venn Diagrams

Using  $X_1$  and  $X_2$  to predict  $Y$ :



Overview

Statistical Control

● Visualizing  
Control

● Example Data Set

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation


Multiple Partial  
Correlation

Wrapping Up



# Venn Diagrams

Using  $X_1$  to predict  $Y$ , while controlling for the effects of  $X_2$ :



Overview

Statistical Control

● Visualizing  
Control

● Example Data Set

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Today's Example Data Set

From Weisberg (1985, p. 240).

“Property taxes on a house are supposedly dependent on the current market value of the house. Since houses actually sell only rarely, the sale price of each house must be estimated every year when property taxes are set. Regression methods are sometimes used to make up a prediction function.”

Overview

Statistical Control

● Visualizing  
Control

● **Example Data Set**

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Erie, Pennsylvania

We have data for 27 houses sold in the mid 1970's in Erie, Pennsylvania:

- $X_1$ : Current taxes (local, school, and county)  $\div$  100 (dollars).
- $X_2$ : Number of bathrooms.
- $X_3$ : Lot size  $\div$  1000 (square feet).
- $X_4$ : Living space  $\div$  1000 (square feet).
- $X_5$ : Number of garage spaces.
- $X_6$ : Number of rooms.
- $X_7$ : Number of bedrooms.
- $X_8$ : Age of house (years).
- $X_9$ : Number of fireplaces.
- $Y$ : Actual sale price  $\div$  1000 (dollars).

Overview

Statistical Control

● Visualizing  
Control

● **Example Data Set**

Partial Correlation


Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up

# Erie, Pennsylvania



Lake Erie

## ***Erie, Pennsylvania***



Jack

## ***Erie, Pennsylvania***



Paula



# Erie, Pennsylvania

For our example today, we will limit the size of the data to a smaller set of variables:

- $X_1$ : Current taxes (local, school, and county)  $\div$  100 (dollars).
- $X_2$ : Number of bathrooms.
- $X_3$ : Living space  $\div$  1000 (square feet).
- $X_4$ : Age of house (years).
- $Y$ : Actual sale price  $\div$  1000 (dollars).

Overview

Statistical Control

● Visualizing  
Control

● Example Data Set

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Correlation Matrix

Variable	$X_1$	$X_2$	$X_3$	$X_4$	$Y$
$X_1$	1.000				
$X_2$	0.876	1.000			
$X_3$	0.832	0.901	1.000		
$X_4$	-0.371	-0.211	-0.178	1.000	
$Y$	0.915	0.924	0.929	-0.310	1.000

- $X_1$ : Current taxes (local, school, and county)  $\div$  100 (dollars).
- $X_2$ : Number of bathrooms.
- $X_3$ : Living space  $\div$  1000 (square feet).
- $X_4$ : Age of house (years).
- $Y$ : Actual sale price  $\div$  1000 (dollars).

Overview

Statistical Control

● Visualizing  
Control

● Example Data Set

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation


Wrapping Up



# Partial Correlation

A partial correlation is a correlation between two variables from which the linear relations, or effects, of another variable(s) have been removed.

$$r_{12.3} = \frac{r_{12} - r_{13}r_{23}}{\sqrt{1 - r_{13}^2}\sqrt{1 - r_{23}^2}}$$



Overview

Statistical Control

Partial Correlation

● Partial Correlation

- Higher-Order Partial Correlation
- Partial Correlation From Regression
- Higher Orders from Regression
- More SPSS

Measurement Errors

Semipartial Correlation

Multiple Partial Correlation

Wrapping Up



# Partial Correlation in SPSS

## Overview

## Statistical Control

## Partial Correlation

### ● Partial Correlation

- Higher-Order Partial Correlation
- Partial Correlation From Regression
- Higher Orders from Regression
- More SPSS

## Measurement Errors

## Semipartial Correlation

## Multiple Partial Correlation

## Wrapping Up

- To perform a partial correlation in SPSS go to Analyze...Correlation...Partial.
- In the Variables box, put the variables you would like to get partial correlations for.
- In the Controlling For box, put the variables you would like to control for.

Partial correlation of  $Y(1)$  (actual sale price ÷ \$1000) with  $X_3(2)$  (living space), controlling for  $X_1(3)$  (current taxes):


$$r_{yx_3 \cdot x_1} = \frac{r_{yx_3} - r_{yx_1}r_{x_3x_1}}{\sqrt{1 - r_{yx_1}^2} \sqrt{1 - r_{x_3x_1}^2}} =$$
$$\frac{0.929 - (0.915 \times 0.832)}{\sqrt{1 - 0.915^2} \sqrt{1 - 0.832^2}} = 0.748$$



# Higher-Order Partial Correlation

The second-order partial correlation is the correlation between two variables with the effects of two other variables being removed:

$$r_{12.34} = \frac{r_{12.3} - r_{14.3}r_{24.3}}{\sqrt{1 - r_{14.3}^2} \sqrt{1 - r_{24.3}^2}}$$



Overview

Statistical Control

Partial Correlation

● Partial Correlation

● **Higher-Order  
Partial Correlation**

● Partial Correlation  
From Regression

● Higher Orders  
from Regression

● More SPSS

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Second-order Partial Correlation

- To perform a partial correlation in SPSS go to Analyze...Correlation...Partial.
- In the Variables box, put the variables you would like to get partial correlations for.
- In the Controlling For box, put the variables you would like to control for - if more than one, a higher-order partial is given.

Overview

Statistical Control

Partial Correlation

● Partial Correlation

● **Higher-Order  
Partial Correlation**

● Partial Correlation  
From Regression

● Higher Orders  
from Regression

● More SPSS

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Second-order Partial Correlation

Partial correlation of  $Y(1)$  (actual sale price  $\div$  \$1000) with  $X_3(2)$  (living space), controlling for  $X_1(3)$  (current taxes) and  $X_2(4)$  (number of bathrooms):

$$r_{12.3} = r_{yx_3.x_1} = 0.748$$

$$r_{14.3} = r_{yx_2.x_1} = 0.628$$

$$r_{24.3} = r_{x_3x_2.x_1} = 0.641$$

$$r_{12.34} = \frac{r_{12.3} - r_{14.3}r_{24.3}}{\sqrt{1 - r_{14.3}^2}\sqrt{1 - r_{24.3}^2}} = \frac{0.748 - (0.628 \times 0.641)}{\sqrt{1 - 0.628^2}\sqrt{1 - 0.641^2}} = 0.578$$

Overview

Statistical Control

Partial Correlation

● Partial Correlation

● Higher-Order  
Partial Correlation

● Partial Correlation  
From Regression

● Higher Orders  
from Regression

● More SPSS

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Partial Correlations from Multiple Correlation

As you could probably guess, partial correlations can be found from regression analyses...

Squared partial correlation:

$$r_{12.3}^2 = \frac{R_{1.23}^2 - R_{1.3}^2}{1 - R_{1.3}^2}$$

- $R_{1.23}^2$  is the  $R^2$  from a multiple regression with 1 being  $Y$  and 2 and 3 being the predictor variables.
- $R_{1.3}^2$  is the  $R^2$  from a simple regression with 1 being  $Y$  and 2 being  $X$  - the single predictor variable.

Overview

Statistical Control

Partial Correlation

● Partial Correlation

● Higher-Order

Partial Correlation

● **Partial Correlation  
From Regression**

● Higher Orders  
from Regression

● More SPSS

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Partial Correlations from Multiple Correlation

Partial correlation of  $Y(1)$  (actual sale price  $\div$  \$1000) with  $X_3(2)$  (living space), controlling for  $X_1(3)$  (current taxes):

$$R_{1.23}^2 = 0.928$$

$$R_{1.3}^2 = 0.838$$

$$r_{12.3}^2 = \frac{R_{1.23}^2 - R_{1.3}^2}{1 - R_{1.3}^2} = \frac{0.928 - 0.838}{1 - 0.838} = 0.865$$

$$r_{12.3} = \sqrt{r_{12.3}^2} = \sqrt{0.865} = 0.748$$

Overview

Statistical Control

Partial Correlation

● Partial Correlation

● Higher-Order

Partial Correlation

● **Partial Correlation  
From Regression**

● Higher Orders  
from Regression

● More SPSS

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Partial Correlations from Multiple Correlation

Alternatively:

$$r_{12.3}^2 = \frac{R_{2.13}^2 - R_{2.3}^2}{1 - R_{2.3}^2}$$

Partial correlation of  $Y(1)$  (actual sale price  $\div$  \$1000) with  $X_3(2)$  (living space), controlling for  $X_1(3)$  (current taxes):

$$R_{2.13}^2 = 0.865$$

$$R_{2.3}^2 = 0.693$$

$$r_{12.3}^2 = \frac{R_{2.13}^2 - R_{2.3}^2}{1 - R_{2.3}^2} = \frac{0.865 - 0.693}{1 - 0.693} = 0.865$$

$$r_{12.3} = \sqrt{r_{12.3}^2} = \sqrt{0.865} = 0.748$$

Overview

Statistical Control

Partial Correlation

- Partial Correlation
- Higher-Order Partial Correlation
- **Partial Correlation From Regression**

- Higher Orders from Regression
- More SPSS

Measurement Errors

Semipartial Correlation

Multiple Partial Correlation

Wrapping Up



# Higher-Order Partial Correlations

Second-order squared partial correlation:

$$r_{12.34}^2 = \frac{R_{1.234}^2 - R_{1.34}^2}{1 - R_{1.34}^2}$$

$$r_{12.34}^2 = \frac{R_{2.134}^2 - R_{2.34}^2}{1 - R_{2.34}^2}$$

Third-order squared partial correlation:

$$r_{12.345}^2 = \frac{R_{1.2345}^2 - R_{1.345}^2}{1 - R_{1.345}^2}$$

$$r_{12.345}^2 = \frac{R_{2.1345}^2 - R_{2.345}^2}{1 - R_{2.345}^2}$$

Overview

Statistical Control

Partial Correlation

● Partial Correlation

● Higher-Order

Partial Correlation

● Partial Correlation

From Regression

● **Higher Orders  
from Regression**

● More SPSS

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Higher-Order Partial Correlations

Partial correlation of  $Y(1)$  (actual sale price  $\div$  \$1000) with  $X_3(2)$  (living space), controlling for  $X_1(3)$  (current taxes) and  $X_2(4)$  (number of bathrooms):

$$R_{1.234}^2 = 0.935$$

$$R_{1.34}^2 = 0.902$$

$$r_{12.34}^2 = \frac{R_{1.234}^2 - R_{1.34}^2}{1 - R_{1.34}^2} = \frac{0.935 - 0.902}{1 - 0.902} = 0.337$$

$$r_{12.34} = \sqrt{r_{12.34}^2} = \sqrt{0.337} = 0.578$$

Overview

Statistical Control

Partial Correlation

● Partial Correlation

● Higher-Order

Partial Correlation

● Partial Correlation

From Regression

● **Higher Orders  
from Regression**

● More SPSS

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Partial Correlations in Regression

You can obtain partial correlations (and semipartial correlations - called part correlations by SPSS) from the regression analysis subroutine:

- Go to Analyze...Regression...Linear.
- Click on Statistics box.
- Check Part and Partial Correlations.

Overview

Statistical Control

Partial Correlation

- Partial Correlation
- Higher-Order Partial Correlation
- Partial Correlation From Regression
- Higher Orders from Regression

● More SPSS

Measurement Errors

Semipartial Correlation

Multiple Partial Correlation

Wrapping Up



# Correction For Attenuation

- Often in behavioral research, variables are not measured without error (non-perfect reliability).
- For these times, the correlation between two variables is dampened by extra noise associated with measurement.
- To correct for this noise, a so-called correction for attenuation accounts for the imperfections in measurement.
- Using these corrected values, one can then build corrected partial correlations.

Correction for attenuation:

$$r_{12}^* = \frac{r_{12}}{\sqrt{r_{11}}\sqrt{r_{22}}}$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

● Correction For  
Attenuation

● Corrected Partial  
Correlation

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Corrected Partial Correlation

Recall the formula for the first-order partial correlation:

$$r_{12.3} = \frac{r_{12} - r_{13}r_{23}}{\sqrt{1 - r_{13}^2}\sqrt{1 - r_{23}^2}}$$

First-order partial correlation correction for attenuation:

$$r_{12.3}^* = \frac{r_{33}r_{12} - r_{13}r_{23}}{\sqrt{r_{11}r_{33} - r_{13}^2}\sqrt{r_{22}r_{33} - r_{23}^2}}$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

● Correction For  
Attenuation

● **Corrected Partial  
Correlation**

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up



# Semipartial Correlation

- Semipartial correlation removes the effects of additional variables from one of the variables under study (typically  $X$ ).

First-order semipartial correlation:

$$r_{1(2.3)} = \frac{r_{12} - r_{13}r_{23}}{\sqrt{1 - r_{23}^2}}$$

Or...

$$r_{2(1.3)} = \frac{r_{12} - r_{13}r_{23}}{\sqrt{1 - r_{13}^2}}$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

**Semipartial  
Correlation**

- Semipartial Correlation from Regression
- Tests of Significance
- Multiple Regression

Multiple Partial  
Correlation

Wrapping Up



# Semipartial Correlation from Regression

First-order semipartial correlation:

$$r_{1(2.3)}^2 = R_{1.23}^2 - R_{1.3}^2$$

Second-order semipartial correlation:

$$r_{1(2.34)}^2 = R_{1.234}^2 - R_{1.34}^2$$

Third-order semipartial:

$$r_{1(2.345)}^2 = R_{1.2345}^2 - R_{1.345}^2$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

● Semipartial  
Correlation from  
Regression

● Tests of  
Significance  
● Multiple  
Regression

Multiple Partial  
Correlation

Wrapping Up



# Semipartial Correlation from Regression

Semipartial correlation of  $Y(1)$  (actual sale price  $\div$  \$1000) with  $X_3(2)$  (living space), controlling  $X_3(2)$  (living space) for  $X_1(3)$  (current taxes):

$$R_{1.23}^2 = 0.928$$

$$R_{1.3}^2 = 0.838$$

$$r_{1(2.3)}^2 = R_{1.23}^2 - R_{1.3}^2 = 0.928 - 0.838 = 0.090$$

$$r_{1(2.3)} = \sqrt{r_{1(2.3)}^2} = \sqrt{0.090} = 0.300$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

● Semipartial  
Correlation from  
Regression

● Tests of  
Significance  
● Multiple  
Regression

Multiple Partial  
Correlation

Wrapping Up



# Semipartial Correlation from Regression

Semipartial correlation of  $Y(1)$  (actual sale price  $\div$  \$1000) with  $X_3(2)$  (living space), controlling  $X_3(2)$  (living space) for  $X_1(3)$  (current taxes) and  $X_2(4)$  (number of bathrooms):

$$R_{1.234}^2 = 0.935$$

$$R_{1.34}^2 = 0.902$$

$$r_{1(2.34)}^2 = R_{1.234}^2 - R_{1.34}^2 = 0.935 - 0.902 = 0.033$$

$$r_{1(2.3)} = \sqrt{r_{1(2.3)}^2} = \sqrt{0.033} = 0.182$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

● Semipartial  
Correlation from  
Regression

● Tests of  
Significance  
● Multiple  
Regression

Multiple Partial  
Correlation

Wrapping Up



# Tests of Significance

Semipartial correlations are found using the formulas developed for the test of the increment in the proportion of variance accounted for in regression:

$$F = \frac{R_{y.12\dots k_1}^2 - R_{y.12\dots k_2}^2 / (k_1 - k_2)}{1 - R_{y.12\dots k_1}^2 / (N - k_1 - 1)},$$

where  $k_1 > k_2$ , with:

- $df_{num} = k_1 - k_2$
- $df_{denom} = N - k_1 - 1$

Overview

Statistical Control

Partial Correlation

Measurement Errors

Semipartial Correlation

● Semipartial Correlation from Regression

● Tests of Significance

● Multiple Regression

Multiple Partial Correlation

Wrapping Up



# Multiple Regression

You can begin to see how multiple regression uses the statistical concepts from today's to control the prediction of the  $Y$  variable:

$$R_{y.1234}^2 = r_{y1}^2 + r_{y(2.1)}^2 + r_{y(3.12)}^2 + r_{y(3.123)}^2$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

● Semipartial  
Correlation from  
Regression

● Tests of  
Significance

● Multiple  
Regression

Multiple Partial  
Correlation

Wrapping Up



# Multiple Partial Correlation

Multiple partial correlation:

$$R_{1.23(4)}^2 = \frac{R_{1.234}^2 - R_{1.4}^2}{\sqrt{1 - R_{1.4}^2}}$$

Two control variables:

$$R_{1.23(45)}^2 = \frac{R_{1.2345}^2 - R_{1.45}^2}{\sqrt{1 - R_{1.45}^2}}$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

● Multiple Partial  
Correlation

● Multiple  
Semipartial  
Correlation

Wrapping Up



# Multiple Semipartial Correlation

Multiple semipartial correlation:

$$R_{1(23.4)}^2 = R_{1.234}^2 - R_{1.4}^2$$

Two control variables:

$$R_{1(23.45)}^2 = R_{1.2345}^2 - R_{1.45}^2$$

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

● Multiple Partial  
Correlation

● Multiple  
Semipartial  
Correlation

Wrapping Up



# Final Thought

- Often times, experimental control is not possible.
- Using statistical techniques, one can get an idea of how variables are related to one another by removing the effects of spurious variables
- Partial and semipartial are statistical techniques to provide post-hoc control.



Overview

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up

● Final Thought

● Next Class



# Next Time

- Midterm handed out.
- Midterm discussion.

Overview

Statistical Control

Partial Correlation

Measurement  
Errors

Semipartial  
Correlation

Multiple Partial  
Correlation

Wrapping Up

● Final Thought

● Next Class